



CHOOSE TREMFYA®, GET TREMFYA®

INDICATIONS

TREMFYA® is indicated for the treatment of adults with moderate to severe plaque psoriasis who are candidates for systemic therapy or phototherapy.

TREMFYA® is indicated for the treatment of adults with active psoriatic arthritis.

SELECTED IMPORTANT SAFETY INFORMATION

TREMFYA® is contraindicated in patients with a history of serious hypersensitivity reaction to guselkumab or to any of the excipients. Serious hypersensitivity reactions, including anaphylaxis, have been reported. TREMFYA® may increase the risk of infection. Instruct patients to seek medical advice if signs or symptoms of clinically important chronic or acute infection occur. If a clinically important or serious infection develops, discontinue TREMFYA® until infection resolves. Evaluate for tuberculosis before treating with TREMFYA®. Avoid use of live vaccines in patients treated with TREMFYA®. Please see related and other [Important Safety Information](#) on last page.



RESOURCES TO HELP YOUR PATIENTS



We are dedicated to helping your patients get started with the TREMFYA® treatment you prescribed. Enrollment starts with you completing

1 FORM

Please see the [Important Safety Information](#) on the last page and the full [Prescribing Information](#). Provide the [Medication Guide](#) to your patients and encourage discussion.

SUPPORT FOR YOUR PATIENTS



In addition, in the midst of current uncertainties surrounding insurance coverage, we want to help your patients understand their coverage options and where Janssen programs may be able to help. Enrollment starts with your patient accepting

**1 PHONE CALL
866-889-5660**

Scan to add the Wegmans Specialty Pharmacy number to your phone



START WITH TREMFYA® SO SIMPLE TRIAL PROGRAM

The TREMFYA® So Simple Trial Program provides patients their first dose of TREMFYA® so they may determine with their provider if it is right for them.

1 **FORM** for easy enrollment

\$0 **COST** to patient

3 Shipment can be authorized within **3 BUSINESS DAYS** after prescription.*

This trial program is open to patients who have commercial insurance, government coverage, or no insurance coverage. However, there is no guarantee of continuous accessibility after the program ends. This program is for medication only. Terms expire at the end of each calendar year and may change.

*Shipment can be authorized within 3 business days of submitting a prescription, pending patient program opt-in and scheduling shipment.

This program is for medication only. Terms expire at the end of each program year and may change.

SUBMIT A COMPLETED PRESCRIPTION ENROLLMENT FORM
(VIA JANSSEN CAREPATH)

Please see the [Important Safety Information](#) on the last page and the full [Prescribing Information](#). Provide the [Medication Guide](#) to your patients and encourage discussion.

FOR COMMERCIALLY
INSURED PATIENTS

STAY ON TREMFYA®

SUBMIT A COMPLETED PRIOR AUTHORIZATION FORM
to the patient's insurance company (via Janssen CarePath or the insurance company directly)

JANSSEN CAREPATH SAVINGS PROGRAM

INSURANCE
COVERAGE
APPROVED

Eligible patients pay **\$5** per injection

with a \$20,000 maximum program benefit per calendar year
See full program requirements at Tremfya.JanssenCarePathSavings.com.

COVERAGE
DELAYED >5
BUSINESS DAYS
OR DENIED

Janssen Link

Patients will receive TREMFYA® **at no cost** until they receive insurance coverage approval

See full program requirements at JanssenCarePath.com/Tremfya/Janssen-Link.

Both programs are unavailable to individuals who use any state or federal government-funded healthcare program to cover a portion of medication costs, such as Medicare, Medicaid, TRICARE, Department of Defense, or Veterans Administration.

These programs are for medication only. Terms expire at the end of each program year and may change.

PATIENTS
NEEDING
EXTRA
SUPPORT

JANSSEN CAREPATH

can provide information about other resources that may be able to help your patients with their out-of-pocket costs

Visit JanssenPrescriptionAssistance.com for more information about affordability programs that may be available.

For more information about Janssen CarePath,
talk with your Janssen Representative.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

TREMFYA® (guselkumab) is contraindicated in patients with a history of serious hypersensitivity reaction to guselkumab or to any of the excipients.

WARNINGS AND PRECAUTIONS

Hypersensitivity Reactions

Serious hypersensitivity reactions, including anaphylaxis, have been reported with postmarket use of TREMFYA®.

Some cases required hospitalization. If a serious hypersensitivity reaction occurs, discontinue TREMFYA® and initiate appropriate therapy.

Infections

TREMFYA® may increase the risk of infection.

Treatment with TREMFYA® should not be initiated in patients with a clinically important active infection until the infection resolves or is adequately treated.

Consider the risks and benefits of treatment prior to prescribing TREMFYA® in patients with a chronic infection or a history of recurrent infection. Instruct patients receiving TREMFYA® to seek medical help if signs or symptoms of clinically important chronic or acute infection occur. If a patient develops a clinically important or serious infection, or is not responding to standard therapy, closely monitor and discontinue TREMFYA® until the infection resolves.

Pre-Treatment Evaluation for Tuberculosis (TB)

Evaluate patients for TB infection prior to initiating treatment with TREMFYA®. Initiate treatment of latent TB prior to administering TREMFYA®. Monitor patients for signs and symptoms of active TB during and after TREMFYA® treatment. Do not administer TREMFYA® to patients with active TB infection.

Immunizations

Prior to initiating TREMFYA®, consider completion of all age-appropriate immunizations according to current immunization guidelines. Avoid use of live vaccines in patients treated with TREMFYA®.

ADVERSE REACTIONS

Most common ($\geq 1\%$) adverse reactions associated with TREMFYA® include upper respiratory infections, headache, injection site reactions, arthralgia, bronchitis, diarrhea, gastroenteritis, tinea infections, and herpes simplex infections.

The overall safety profile observed in patients with psoriatic arthritis is generally consistent with the safety profile in patients with plaque psoriasis, with the addition of bronchitis and neutrophil count decreased.

Please read the full Prescribing Information and Medication Guide for TREMFYA®. Provide the Medication Guide to your patients and encourage discussion.

cp-82625v3

